

A Case for Customizing Estimators: Coordinated Samples

Edith Cohen
Microsoft Research
Mountain View, CA, USA
edith@cohenwang.com

Haim Kaplan
School of Computer Science
Tel Aviv University, Israel
haimk@cs.tau.ac.il

ABSTRACT

Random samples are a popular method of summarizing data and allow queries posed over the data to be approximated by applying an appropriate estimator to the sample.

Performance on a particular query, however, hinges on estimator selection. The choice of estimators is subjected to global requirements, such as unbiasedness and range restrictions on the estimate value, and ideally, we seek estimators that are both efficient to derive and apply and optimal in that they are not dominated by other estimators. Nevertheless, for a given data domain, sampling scheme, and query, there are many applicable estimators. In this work, we aim to understand and leverage this choice. When the data exhibits patterns that we can learn or observe, we can *customize* estimators to these patterns to obtain better performance while maintaining certain guarantees for all possible data in the domain.

Through a case study of coordinated sampling and focusing on estimators that are variance optimal, nonnegative, and unbiased, we set some foundations for the derivation of customized estimators and demonstrate its usefulness for data analysis applications.

1. INTRODUCTION

Random samples are extensively used for scalable analysis of massive data. The samples facilitate approximate processing of queries posed over the original data, when exact processing is too resource consuming or when the original data is no longer available. Random samples have a distinct advantages over other synopsis in their flexibility, in terms of supported queries.

Queries performed over the data range from basic subset statistics, such as sums, moments, and averages, and more complex relations: distinct counts, size of set intersections, and difference norms. The value of a sample hinges on the accuracy within which we can estimate query results. In turn, this boils down to the *estimators* we use, which are the functions we apply to the sample to produce the estimate.

We seek estimators that satisfy some *global* properties, which hold for all possible data in our domain. Some global properties we consider are

- *Range restriction* of estimates: since the estimate is often used as a substitute of the true value, we would like it to be from the same range as the query result. Some common restrictions are *nonnegativity* (when the estimated range is), or *boundedness* which means that the range of estimates is bounded by some function of the query result.
- *Unbiasedness*, which means that the expectation of the estimate is equal to the query result. Unbiasedness is particularly important when we estimate a sum aggregate by summing estimates of components, and wish the relative error to decrease with aggregation.
- *Finite variance* (implied by boundedness but less restrictive)

Perhaps the most basic quality measure of an estimator is its variance. The variance, however, depends on the input data set, and in general there is no single estimator that is optimal for all data values in our domain. We therefore aim for *variance optimality* – meaning that strict improvement is not possible without violating some global properties. More precisely, there is no estimator with at most the variance of our estimator on all data and strictly lower variance on some data. We also consider *variance competitiveness* [8] – meaning that the variance on each data is not too far off the best possible for the data subject to the global properties. Variance competitiveness provides “worst case” performance guarantees over data in our domain. It may not always be desirable or achievable but is useful when we need some guarantees for all possible data.

We treat the problem of deriving estimators from an optimization perspective, looking for estimators that satisfy global properties and optimality but at the same time, are tailored to perform better when the data follows some stated patterns. The first component is to understand the set of applicable estimators, those satisfying the global properties we are after and (Pareto) optimality. We then hope to leverage the freedom we have in estimator selection to derive *customized* estimators that performs better on recurring patterns we can learn or observe in the data, such as sparsity of certain relations between entries.

We explore this process for coordinated shared-seed sampling [1, 13, 11, 12, 3, 2, 4, 5, 9] (see [8, 9] for overview).

The sampling scheme is widely applicable and thus the development of good estimators is independently interesting. It is also sufficiently complex to be technically challenging and yet simple enough to allow for a comprehensive understanding. In particular, variance competitiveness is attainable [8] and there is a complete characterization of functions for which estimators with certain global properties exist [8].

Example 1 Dataset with 3 instances and queries

Instances $i \in \{1, 2, 3\}$ and keys $k \in \{a, b, c, d, e, f, g, h\}$:

	a	b	c	d	e	f	g	h
1	0.95	0	0.23	0.70	0.10	0.42	0	0.32
2	0.15	0.44	0	0.80	0.05	0.50	0.20	0
3	0.25	0	0	0.10	0	0.22	0	0

Example queries over selected keys $H \subset [a-h]$. L_p difference, L_p^p , which is the p th power of L_p difference and a sum aggregate which can be used to estimate the L_p difference, L_{p+}^p : asymmetric (increase only) L_p^p , the sum of the increase-only and the decrease-only changes (decrease only is obtained by switching the roles of v_1 and v_2) is L_p^p , but each component is a useful metric for asymmetric change. G an “arbitrary” sum aggregate, illustrating versatility (work applicable to any function for which estimators exist)

$$L_p(H) = (\sum_{k \in H} |v_1^{(k)} - v_2^{(k)}|^p)^{1/p}$$

$$L_p^p(H) = \sum_{k \in H} |v_1^{(k)} - v_2^{(k)}|^p$$

$$L_{p+}^p(H) = \sum_{k \in H} \max\{0, v_1^{(k)} - v_2^{(k)}\}^p$$

$$G(H) = \sum_{k \in H} |v_1^{(k)} - 2v_2^{(k)} + v_3^{(k)}|^2$$

sum aggregate	tuple function
L_p^p	$\text{RG}_p(\mathbf{v}) = (\max(\mathbf{v}) - \min(\mathbf{v}))^p$
L_{p+}^p	$\text{RG}_{p+}(v_1, v_2) = \max\{0, v_1 - v_2\}^p$
G	$g(v_1, v_2, v_3) = v_1 + v_3 - 2v_2 ^2$

$$L_1(\{b, c, e\}) = |0 - 0.44| + |0.23 - 0| + |0.10 - 0.05| = 0.71$$

$$L_2^2(\{c, f, h\}) = (0.23 - 0)^2 + (0.50 - 0.42)^2 + (0.32 - 0)^2 \approx 0.16$$

$$L_2(\{c, f, h\}) = \sqrt{L_2^2(\{3, 6, 8\})} \approx 0.40$$

$$L_{1+}(\{b, c, e\}) = \max\{0, 0 - 0.44\} + \max\{0, 0.23 - 0\} + \max\{0, 0.10 - 0.05\} = 0.235$$

$$G(\{b, d\}) = |0 - 2 * 0.44 + 0|^2 + |0.7 - 2 * 0.8 + 0.1|^2 \approx 1.18$$

Coordinated sampling is particularly useful when data sets have the form of multiple *instances*, which are weight assignments over the same set of *keys*. Different instances may correspond to snapshots, activity logs, measurements, or repeated surveys at different times or locations. Coordination of the samples of different instances allows for more accurate estimates of aggregates such as distinct counts and similarity measures. Such aggregates often either have the form of a sum over keys of a “tuple” function applied to the values of the key in the different instances, or otherwise can be expressed as functions of such sums: Distinct count is a sum aggregate of logical OR and the L_p difference is the

p th root of L_p^p , which sum-aggregates exponentiated range functions $\text{RG}_p(\mathbf{v}) = (\max(\mathbf{v}) - \min(\mathbf{v}))^p$ ($p > 0$). Example 1 illustrates a data set, example aggregate queries, and respective tuple functions.

We focus on estimating “tuple” functions defined over the tuple of the values of a single key. Sum aggregates can be estimated by summing up tuple estimators. Tuple estimates typically have high variance, since most of all of the entries are missing from the sample. We therefore insist on unbiased and (at least) pairwise independence of the tuple estimates, which allow the relative error to decrease with aggregation [8]. Since the tuple functions we are interested in are non-negative, we also require the tuple estimates to be nonnegative [8] (results extend to any one-sided range restriction on the estimates).

The pointwise-optimal range: We express the range of estimators that are variance⁺ optimal, where we define variance⁺ optimality as variance optimality over unbiased nonnegative estimators. The *pointwise optimal range* of estimates (Section 3) is defined for an outcome, conditioned on estimate values on all “less-informative” outcomes, and includes the range of estimate values that are optimal with respect to at least one data vector that is consistent with the outcome. We show that being “in range” almost everywhere is necessary for variance⁺ optimality and sufficient for unbiasedness and nonnegativity, when an unbiased nonnegative estimator exists for the function f .

The L and U estimators: The lower extreme of the pointwise optimal range is the *L estimator* (Section 4), which has a compelling combination of properties. It satisfies both our quality measures, being variance⁺ optimal and 4-competitive. The L estimator is *monotone*, meaning that when fixing the data vector, the estimate value is monotone non-decreasing with the information we can glean from the outcome (set of vectors consistent with our sample). In fact, the L estimator is the *unique variance⁺ optimal monotone estimator* and thus dominates (in terms of variance pointwise) the Horvitz-Thompson estimator [10] (which is also unbiased, nonnegative, and monotone), when the latter is applicable. We show that the competitive ratio of 4 of the L estimator is tight in the sense that there is a family of functions on which the supremum of the ratio, over functions and data vectors, is 4. The ratio can be lower, however, for specific functions. Finally, we give a simple expression for the L estimator which allows it to be efficiently computed by numeric integration or a closed form.

The upper extreme of the pointwise optimal range is the *U estimator* (Section 6), which is unbiased, nonnegative, and has finite variances. We show that under some conditions on f that are satisfied by natural functions including the exponentiated range, the U estimator is also variance⁺ optimal.

Order-based optimality: One notion useful for customization is *order optimality* [6]: An estimator is $\prec^+$ -optimal with respect to some partial order $\prec$ on data vectors if any other (nonnegative unbiased) estimator with lower variance

on some data v must have strictly higher variance on some data that precedes v . Order-based optimality implies variance optimality, but not vice versa. By specifying an order which prioritizes more likely patterns in the data, we can customize the estimator to these patterns.

We show (Section 5) how to construct a $\prec^+$ -optimal non-negative unbiased estimators for any function and order $\prec$ for which the estimator exists. We show that when the data domain is discrete, such estimators always exist whereas continuous domains require some natural convergence properties of $\prec$. Moreover, the L estimator is $\prec^+$ -optimal with respect to $z \prec v \iff f(z) < f(v)$. The U estimator, under some conditions, is $\prec^+$ -optimal with respect to the reverse order.

Practical implications: In [7], we evaluate L_p difference estimators derived as p th roots of sums of our L and U estimators for exponentiated range functions RG_p ($p > 0$). We apply the L_1 and L_2 estimators to samples of data sets with different characteristics: IP flow records exhibited larger differences between bandwidth usage assumed by a flow key (IP source destination pair, port, and protocol) in different times. The surnames dataset (frequencies of surnames in published books in different years) had more similar values. Accordingly, the U estimator, which is optimized for large differences dominated on the IP flow records dataset whereas the L estimator dominated on the surnames dataset. This demonstrates the potential value in selecting a custom estimator.

More generally, the study shows that we obtain accurate estimates even when only a small fraction of entries is sampled, using either estimator, and also demonstrate the value of competitiveness of the L estimator: Whereas it can outperform the U estimator by orders of magnitude, it can be outperformed only by a small factor.

2. PRELIMINARIES

We briefly summarize model and concepts on estimator optimality [6] and coordinated shared-seed sampling [8].

Sampling model: The data is a vector $v = (v_1, v_2, \dots, v_r) \in V = V^r$, where V is some subset of the reals. The *sampling scheme* we use is specified by continuous functions $\tau = (\tau_1, \dots, \tau_r)$ on $[0, 1]$ with range containing $(\min V, \max V)$. The output of the sampling is the outcome $S \equiv S(u, v)$, which depends on the data and a random seed $u \in U[0, 1]$. We treat the outcome as a set where the i th entry is included in S if and only if v_i is at least $\tau_i(u)$:

$$i \in S \iff v_i \geq \tau_i(u)$$

but also assume that the seed u is available with it. A special case of particular interest is PPS sampling, when $\tau_i(u)$ are linear functions: There is a fixed vector τ^* such that $\tau_i(u) \equiv u\tau_i^*$.

With each outcome $S(u, v)$, we can identify the set $V^*(S)$

of all data vectors that are consistent with it:

$$V^*(S) \equiv V^*(u, v) = \{z \mid \forall i \in [r], i \in S \wedge z_i = v_i \vee i \notin S \wedge z_i < \tau_i(u)\}.$$

The set $V^*(u, v)$ is increasing with u , which means we have less information on the data when u is larger. For two different outcomes S_1 and S_2 , the sets $V^*(S_1)$ and $V^*(S_2)$ must be either disjoint or one is contained in the other.

For any two vectors, the set of u values on which the outcomes $S(u, v)$ and $S(u, z)$ are the same is a suffix of $(0, 1]$ that is open to the left:

$$\begin{aligned} \forall \rho \in (0, 1] \forall v, \\ z \in V^*(\rho, v) \implies \exists \epsilon > 0, \forall x \in (\rho - \epsilon, 1], z \in V^*(x, v) \end{aligned} \quad (1)$$

Example 2 Coordinated PPS sampling for Example 1

Consider shared-seed coordinated sampling, where each of the instances A,B,C is PPS sampled with threshold $\tau^* = 1$. In this particular case, each entry is sampled with probability equal to its value. To coordinate the samples, we draw $u^{(k)} \in U[0, 1]$, independently for different keys. A key k is sampled in instance i if and only if $v_i^{(k)} \geq u^{(k)}$. $V^*(S^{(k)})$ contains all vectors consistent with the sampled entries and with value at most $u^{(k)}$ in unsampled entries.

key	a	b	c	d	e	f	g	h
1	0.95	0	0.23	0.70	0.10	0.42	0	0.32
2	0.15	0.44	0	0.80	0.05	0.50	0.20	0
3	0.25	0	0	0.10	0	0.22	0	0
$u^{(k)}$	0.32	0.21	0.04	0.23	0.84	0.70	0.15	0.64

The outcomes for the different keys are: $S^{(a)} = (0.95, *, *)$, $S^{(b)} = (*, 0.44, *)$, $S^{(c)} = (0.23, *, *)$, $S^{(d)} = (0.7, 0.8, *)$, $S^{(e)} = S^{(f)} = S^{(h)} = (*, *, *)$, $S^{(g)} = (*, 0.2, *)$. The sets of vectors consistent with the outcomes are $V^*(S^{(a)}) = \{0.95\} \times [0, 0.32]^2$ and $V^*(S^{(h)}) = [0, 0.64]^3$.

Estimators: We are interested in estimating, from the outcome, a function $f : V$ which maps V to the nonnegative reals. We apply an *estimator* $\hat{f}$ to the outcome (including the seed) and use the notation $\hat{f}(u, v) \equiv \hat{f}(S(u, v))$. When the domain is continuous, we assume $\hat{f}$ is (Lebesgue) integrable.

Two estimators $\hat{f}_1$ and $\hat{f}_2$ are *equivalent* if for all data v , $\hat{f}_1(u, v) = \hat{f}_2(u, v)$ with probability 1, which is the same as

$$\hat{f}_1 \text{ and } \hat{f}_2 \text{ are equivalent} \iff \forall v \forall \rho \in (0, 1], \quad (2)$$

$$\lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}_1(u, v) du}{\rho - \eta} = \lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}_2(u, v) du}{\rho - \eta}.$$

An estimator $\hat{f}$ is *nonnegative* if $\forall S, \hat{f}(S) \geq 0$ and is *unbiased* if $\forall v, E[\hat{f}|v] = f(v)$. An estimator has *finite variance* on v if $\int_0^1 \hat{f}(u, v)^2 du < \infty$ (the expectation of the square is finite) and is *bounded* on v if $\sup_{u \in (0, 1]} \hat{f}(u, v) < \infty$. If a nonnegative estimator is bounded on v , it also has finite variance for v . An estimator is *monotone* on v if when fixing v and considering outcomes consistent with v , the estimate value is non decreasing with the information on the

data that we can glean from the outcome, that is, $\hat{f}(u, v)$ is non-increasing with u . We say that an estimator is bounded, has finite variances, or is monotone, if the respective property holds for all $v \in \mathbf{V}$.

The lower bound function. For $Z \subset \mathbf{V}$, we define $\underline{f}(Z) = \inf\{f(v) \mid v \in Z\}$ as the infimum of f on Z . We use the notation $\underline{f}(S) \equiv \underline{f}(V^*(S))$, $\underline{f}(\rho, v) \equiv \underline{f}(V^*(\rho, v))$. When v is fixed, we use $\underline{f}^{(v)}(u) \equiv \underline{f}(u, v)$. Some properties which we need in the sequel are [8]:

$$\bullet \forall v, \underline{f}^{(v)}(u) \text{ is} \quad (3)$$

montone non increasing and left-continuous.

$$\bullet \hat{f} \text{ is unbiased and nonnegative} \implies \quad (4)$$

$$\forall v, \forall \rho, \int_{\rho}^1 \hat{f}(u, v) du \leq \underline{f}^{(v)}(\rho). \quad (5)$$

The lower bound function $\underline{f}^{(v)}$, and its lower hull $H_f^{(v)}$, are instrumental in capturing existence of estimators with desirable properties [8]:

$$\bullet \exists \text{ unbiased nonnegative } f \text{ estimator} \iff \quad (6)$$

$$\forall v \in \mathbf{V}, \lim_{u \rightarrow 0^+} \underline{f}^{(v)}(u) = f(v). \quad (7)$$

• If f satisfies (7),

$\exists$ unbiased nonnegative estimator with finite variance for v

$$\iff \int_0^1 \left(\frac{dH_f^{(v)}(u)}{du} \right)^2 du < \infty. \quad (8)$$

$\exists$ unbiased nonnegative estimator that is bounded on v

$$\iff \lim_{u \rightarrow 0^+} \frac{f(v) - \underline{f}^{(v)}(u)}{u} < \infty. \quad (9)$$

Example 3 illustrates the lower bound functions and respective lower hull for RG_{p+} .

Partially specified estimators. We use *partial specifications* $\hat{f}$ of (nonnegative and unbiased) estimators, which are specified on a set of outcomes $\mathcal{S}$ so that

$$\forall v \exists \rho_v \in [0, 1], S(u, v) \in \mathcal{S} \text{ a.e. for } u > \rho_v \wedge \\ S(u, v) \notin \mathcal{S} \text{ a.e. for } u \leq \rho_v.$$

When $\rho_v = 0$, we say that the estimator is *fully specified* for v . We also require that $\hat{f}$ is nonnegative where specified and satisfies

$$\forall v, \rho_v > 0 \implies \int_{\rho_v}^1 \hat{f}(u, v) du \leq f(v) \quad (10a)$$

$$\forall v, \rho_v = 0 \implies \int_{\rho_v}^1 \hat{f}(u, v) du = f(v). \quad (10b)$$

LEMMA 2.1. *If f satisfies (7) (has a nonnegative unbiased estimator), then any partially specified estimator can be extended to an unbiased nonnegative estimator.*

v -optimal extensions and estimators. Given a partially specified estimator $\hat{f}$ so that $\rho_v > 0$ and $M = \int_{\rho_v}^1 \hat{f}(u, v) du$,

a *v -optimal extension* is an extension which is fully specified for v and minimizes variance for v (amongst all such extensions). The *v -optimal extension* is defined on outcomes $S(u, v)$ for $u \in (0, \rho_v]$ and satisfies

$$\begin{aligned} \min \int_0^{\rho_v} \hat{f}(u, v)^2 du \\ \text{s.t. } \int_0^{\rho_v} \hat{f}(u, v) du = f(v) - M \\ \forall u, \int_u^{\rho_v} \hat{f}(x, v) dx \leq \underline{f}^{(v)}(u) - M \\ \forall u, \hat{f}(u, v) \geq 0 \end{aligned} \quad (11)$$

For $\rho_v \in (0, 1]$ and $M \in [0, \underline{f}^{(v)}(\rho_v)]$, we define the function $\hat{f}^{(v, \rho_v, M)} : (0, \rho_v] \rightarrow \mathbb{R}_+$ as the solution of

$$\hat{f}^{(v, \rho_v, M)}(u) = \inf_{0 \leq \eta < u} \frac{\underline{f}^{(v)}(\eta) - M - \int_u^{\rho_v} \hat{f}^{(v, \rho_v, M)}(u) du}{\rho - \eta}. \quad (12)$$

Geometrically, the function $\hat{f}^{(v, \rho_v, M)}$ is the negated derivative of the lower hull of the lower bound function $\underline{f}^{(v)}$ on $(0, \rho_v)$ and the point (ρ_v, M) .

THEOREM 2.1. *Given a partially specified estimator $\hat{f}$ so that $\rho_v > 0$ and $M = \int_{\rho_v}^1 \hat{f}(u, v) du$, then $\hat{f}^{(v, \rho_v, M)}$ is the unique (up to equivalence) v -optimal extension of $\hat{f}$.*

The *v -optimal* estimates are the minimum variance extension of the empty specification. We use $\rho_v = 1$ and $M = 0$ and obtain $\hat{f}^{(v)} \equiv \hat{f}^{(v, 1, 0)}$. $\hat{f}^{(v)}$ is the solution of

$$\hat{f}^{(v)}(u) = \inf_{0 \leq \eta < u} \frac{\underline{f}^{(v)}(\eta) - \int_u^1 \hat{f}^{(v)}(u) du}{\rho - \eta}, \quad (13)$$

which is the negated slope of the lower hull of the lower bound function $\underline{f}^{(v)}$. This is illustrated in Example 3.

Variance⁺ and order-based optimality. An estimator is *variance⁺-optimal* if there is no nonnegative unbiased estimator with same or lower variance on all data and strictly lower on some data. We also consider order-based optimality with respect to a partial order $\prec$ on $\mathbf{V}$: An estimator $\hat{f}$ is $\prec^+$ -optimal if there is no other nonnegative unbiased estimator with strictly lower variance on some data v and at most the variance of $\hat{f}$ on all vectors that precede v . Order-based optimality (with respect to some $\prec$) implies variance optimality but the converse is generally not true [6].

Variance competitiveness An estimator $\hat{f}$ is *c-competitive* if

$$\forall v, \int_0^1 \left(\hat{f}(u, v) \right)^2 du \leq c \inf_{\hat{f}'} \int_0^1 \left(\hat{f}'(u, v) \right)^2 du,$$


where the infimum is over all unbiased nonnegative estimators of f . When the estimator is unbiased, the expectation of the square is closely related to variance, and an estimator

Example 3 Lower bound function and its lower hull

Consider $\text{RG}_{p+}(v_1, v_2) = \max\{0, v_1 - v_2\}^p$ (see Example 1) over the domain $\mathbf{V} = [0, 1]^2$ and PPS sampling with $\tau_1^* = \tau_2^* = 1$ (as in Example 2). The lower bound function for data $\mathbf{v} = (v_1, v_2)$ is

$$\text{RG}_{p+}(u, \mathbf{v}) = \max\{0, v_1 - \max\{v_2, u\}\}^p.$$

The figures below illustrate $\text{RG}_{p+}^{(\mathbf{v})}(u)$ and its lower hull for the data vectors $(0.6, 0.2)$ and $(0.6, 0)$ and $p = \{0.5, 1, 2\}$. For $u > 0.2$, the outcome when sampling both vectors is the same, and thus the lower bound function is the same. For $u \leq 0.2$, the outcomes diverge. For $p \leq 1$, $\text{RG}_{p+}^{(\mathbf{v})}(u)$ is concave and the lower hull is linear on $(0, v_1]$. For $p > 1$, the lower hull coincides with $\text{RG}_{p+}^{(\mathbf{v})}(u)$ on some interval $(a, v_1]$ and is linear on $(0, a]$. When $v_2 = 0$, $\text{RG}_{p+}^{(\mathbf{v})}(u)$ is equal to its lower hull.



The $\mathbf{v}$ -optimal estimates are the negated slopes of the lower hulls. They are 0 when $u \in (0.6, 1]$, since these outcomes are consistent with data on which $\text{RG}_{p+} = 0$. They are constant for $u \in (0, v_1]$ when $p \leq 1$. Observe that for $u \in (0.2, 0.6]$, the $\mathbf{v}$ -optimal estimates are different even though the outcome of sampling the two vectors is the same – demonstrates that it is not possible to simultaneously minimize the variance of the two vectors.

that minimizes one also minimizes the other.

$$\text{VAR}[\hat{f}|\mathbf{v}] = \int_0^1 \hat{f}(u, \mathbf{v})^2 du - f(\mathbf{v})^2 \quad (14)$$

3. THE POINTWISE OPTIMAL RANGE

We say that an estimator $\hat{f}$ is $\mathbf{v}$ -optimal at an outcome $S(u, \mathbf{v})$ if it satisfies (13). For an outcome $S(\rho, \mathbf{v})$, we are interested in the *range* of $\mathbf{z}$ -optimal estimates at S for all $\mathbf{z} \in V^*(S)$, with respect to a value M , which captures the contribution to the expectation of the estimator made by outcomes which are less informative than S .

$$\lambda(\rho, \mathbf{v}, M) = \inf_{0 \leq \eta < \rho} \frac{f(\eta, \mathbf{v}) - M}{\rho - \eta} \quad (15)$$

$$\lambda_U(\rho, \mathbf{v}, M) \equiv \lambda_U(S, M) = \sup_{\mathbf{z} \in V^*(\rho, \mathbf{v})} \lambda(\rho, \mathbf{z}, M) \quad (16)$$

$$\begin{aligned} \lambda_L(\rho, \mathbf{v}, M) &\equiv \lambda_L(S, M) = \inf_{\mathbf{z} \in V^*(\rho, \mathbf{v})} \lambda(\rho, \mathbf{v}, M) \\ &= \inf_{\mathbf{z} \in V^*(\rho, \mathbf{v})} \inf_{0 \leq \eta < \rho} \frac{f(\eta, \mathbf{z}) - M}{\rho - \eta} \\ &= \frac{f(\rho, \mathbf{v}) - M}{\rho} \end{aligned} \quad (17)$$

To verify equality (17), observe that from left continuity of $\underline{f}(u, \mathbf{z})$,

$$\inf_{\eta < \rho, \mathbf{z} \in V^*(S)} \underline{f}(\eta, \mathbf{z}) = \underline{f}(\rho, \mathbf{v})$$

and that the denominator $\rho - \eta$ is maximized at $\eta = 0$. $\lambda(\rho, \mathbf{v}, M)$ is the $\mathbf{v}$ -optimal estimate at ρ , given a specifica-




Figure 1: Lower bound functions for vectors $\mathbf{v}, \mathbf{z}, \mathbf{w}$. Outcomes are consistent for all $x \geq u$: $S(x, \mathbf{v}) = S(x, \mathbf{z}) = S(x, \mathbf{w}) \equiv S_x$. The figure illustrates the $\mathbf{y}$ -optimal estimates $\lambda(u, \mathbf{y}, M)$ at u given M for $\mathbf{y} \in \{\mathbf{v}, \mathbf{z}, \mathbf{w}\}$. The estimates are the negated slopes of the lower hull of the point (u, M) and the lower bound function $\underline{f}^{(\mathbf{y})}$. The optimal range at S_u given M is lower-bounded by $\mathbf{w}$, that is $\lambda_L(S_u, M) = \lambda(u, \mathbf{w}, M)$, and upper-bounded by $\mathbf{v}$, $\lambda_U(S_u, M) = \lambda(u, \mathbf{v}, M)$. The figure illustrates the general property that the optimal range is lower bounded by the $\mathbf{w}$ which satisfies $f(\mathbf{w}) = \underline{f}(\mathbf{w}, u)$.

tion of the estimator $\hat{f}(u, v)$ for $u \in (\rho, 1]$ with $\int_{\rho}^1 \hat{f}(u, v) du = M$. In short, we refer to $\lambda(\rho, v, M)$ as the v -optimal estimate at ρ given M . Geometrically, $\lambda(\rho, v, M)$ is the negated slope of the lower hull of $\underline{f}^{(v)}$ and the point (ρ, M) . $\lambda_U(S, M)$ and $\lambda_L(S, M)$, respectively, are the supremum and infimum of the *range* of z -optimal estimates at S given M . Figure 1 illustrates an outcome S and the optimal range at S given M . We can see how the lower endpoint of the range is realized by a vector with f value equal to the lower bound at S , as in equality (17).

When $\hat{f}$ is given for $u \in (\rho, 1]$, we use $M = \int_{\rho}^1 \hat{f}(u, v) du$ and abbreviate the notations by removing M to $\lambda(\rho, v)$, $\lambda_U(S)$, and $\lambda_L(S)$.

We say that the estimator $\hat{f}$ is *in-range* (in the optimal range) at outcome $S(\rho, v)$ if

$$\lambda_L(S) \leq \hat{f}(S) \leq \lambda_U(S). \quad (18)$$

Writing (18) explicitly, we obtain

$$\hat{f}(\rho, v) \geq \lambda_L(\rho, v) = \frac{f(\rho, v) - \int_{\rho}^1 \hat{f}(u, v) du}{\rho} \quad (19a)$$

$$\begin{aligned} \hat{f}(\rho, v) &\leq \lambda_U(\rho, v) \\ &= \sup_{z \in V^*(S)} \inf_{0 \leq \eta < \rho} \frac{f(\eta, z) - \int_{\rho}^1 \hat{f}(u, v) du}{\rho - \eta} \end{aligned} \quad (19b)$$

Two special solutions that we study are the L estimator ($\hat{f}^{(L)}$, see Section 4) and the U estimator ($\hat{f}^{(U)}$, see Section 6), which respectively solve (19a) and (19b) with equalities. For all $\rho \in (0, 1]$ and v , $\hat{f}^{(L)}$ minimizes and $\hat{f}^{(U)}$ maximizes $\int_{\rho}^1 \hat{f}(u, v) du$ among all solutions of (18).

We show that being in-range (satisfying (18) for all outcomes S) is sufficient for nonnegativity and unbiasedness.

LEMMA 3.1. *If f satisfies (7) then any in-range estimator is unbiased and nonnegative.*

PROOF. For nonnegativity, it suffices to show that a solution of (18) satisfies (5), since (19a) and (5) together imply nonnegativity. Assume to the contrary that a solution $\hat{f}$ violates (5) and let ρ be the supremum of x satisfying $\int_x^1 \hat{f}(u, v) du > \underline{f}(x, v)$. From (3), which is monotonicity and left-continuity of $\underline{f}(x, v)$, we have $\int_{\rho}^1 \hat{f}(u, v) du = \underline{f}(\rho, v)$. Since $\int_x^1 \hat{f}(u, v) du$ is continuous in x , and $\underline{f}^{(v)}$ left-continuous, there must be $\delta > 0$ so that

$$\forall x \in [\rho - \delta, \rho), \int_x^1 \hat{f}(u, v) du > \underline{f}(x, v). \quad (20)$$

Let $x \in [\rho - \delta, \rho)$ and $M(x) = \int_x^1 \hat{f}(u, v) du$. From (20),

$M(x) > \underline{f}(x, v)$. We have that

$$\begin{aligned} \hat{f}(x, v) &\leq \sup_{z \in V^*(x, v)} \inf_{0 \leq \eta < x} \frac{\underline{f}(\eta, z) - M(x)}{x - \eta} \\ &\leq \sup_{z \in V^*(x, v)} \inf_{0 \leq \eta < x} \frac{\underline{f}(\eta, z) - \underline{f}(x, v)}{x - \eta} \\ &\leq \sup_{z \in V^*(x, v)} \lim_{\eta \rightarrow x^-} \frac{\underline{f}(\eta, z) - \underline{f}(x, v)}{x - \eta} \\ &= \lim_{\eta \rightarrow x^-} \frac{\underline{f}(\eta, v) - \underline{f}(x, v)}{x - \eta} = -\frac{\partial \underline{f}(x, v)}{\partial x^-} \end{aligned}$$

Since this holds for all $x \in (\rho - \delta, \rho)$, we obtain that $\int_{\rho-\delta}^{\rho} \hat{f}(x, v) dx \leq \underline{f}(\rho - \delta, v) - \underline{f}(\rho, v)$. Therefore, $\int_{\rho-\delta}^1 \hat{f}(x, v) dx \leq \underline{f}(\rho - \delta, v)$, which contradicts (20).

We now establish unbiasedness, using (19a), and $\underline{f}(u, v)$ being non increasing in u , we obtain that $\forall u \forall \rho > u$,

$$\begin{aligned} \hat{f}(u, v) &\geq \frac{f(u, v) - \int_u^1 \hat{f}(x, v) dx}{u} \\ &\geq \frac{f(\rho, v) - \int_u^1 \hat{f}(x, v) dx}{u} \end{aligned} \quad (21)$$

We argue that

$$\forall v \forall \rho > 0, \lim_{x \rightarrow 0} \int_x^1 \hat{f}(u, v) du \geq \underline{f}(\rho, v). \quad (22)$$

To prove (22), define $\Delta(x) = \underline{f}(\rho, v) - \int_x^1 \hat{f}(u, v) du$ for $x \in (0, \rho]$. We show that $\int_{x/2}^x \hat{f}(u, v) du \geq \Delta(x)/4$. To see this, assume to the contrary that $\int_y^x \hat{f}(u, v) du \leq \Delta(x)/4$ for all $y \in [x/2, x]$. Then from (21), the value of $\hat{f}(u, v)$ for $u \in [x/2, x]$ must be at least $(3/4)\Delta(x)/x$. Hence, the integral over the interval $[x/2, x]$ is at least $(3/8)\Delta(x)$ which is a contradiction. We can now apply this iteratively, obtaining that $\Delta(\rho/2^i) \leq (3/4)^i \Delta(\rho)$. Thus, the gap $\Delta(x)$ diminishes as $x \rightarrow 0$ and we established (22).

Since (22) holds for all $\rho \geq 0$, then $\lim_{u \rightarrow 0} \int_u^1 \hat{f}(u, v) du = \lim_{u \rightarrow 0} \underline{f}(u, v) = f(v)$ (using (7)). Combining that we already established (5) we obtain $\lim_{u \rightarrow 0} \int_u^1 \hat{f}(u, v) du = f(v)$. $\square$

We next show that being in-range is necessary for optimality. For our analysis of order-based optimality (Section 5), we need to slightly refine the notion of variance⁺-optimality to be with respect to a partially specified estimator $\hat{f}$ and a subset of data vectors $Z \subset \mathbf{V}$.

An extension of $\hat{f}$ that is fully specified for all vectors in Z is variance⁺-optimal on Z if any other extension with strictly lower variance on at least one $v \in Z$ has a strictly higher variance on at least one $z \in Z$. We say that a partial specification is *in-range with respect to Z* if:

$$\begin{aligned} &\forall v \in Z, \text{ for } \rho \in (0, \rho_v] \text{ almost everywhere,} \\ &\inf_{z \in Z \cap V^*(\rho, v)} \lambda(\rho, z) \leq \hat{f}(\rho, v) \leq \sup_{z \in Z \cap V^*(\rho, v)} \lambda(\rho, z) \end{aligned} \quad (23)$$

Using (2), (23) is the same as requiring that $\forall v \forall \rho \in (0, \rho_v]$, when fixing the estimator on $S(u, v)$ for $u \geq \rho$, then

$$\inf_{z \in Z \cap V^*(\rho, v)} \lambda(\rho, z) \leq \lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} \leq \sup_{z \in Z \cap V^*(\rho, v)} \lambda(\rho, z) \quad (24)$$

We show that a necessary condition for variance⁺-optimality with respect to a partial specification and Z is that almost everywhere, estimates for outcomes consistent with vectors in Z are in-range for Z . Formally:

THEOREM 3.1. *An extension is variance⁺-optimal on Z only if (23) holds.*

PROOF. Consider an (nonnegative unbiased) estimator $\hat{f}$ that violates (23) for some $v \in Z$ and ρ . We show that there is an alternative estimator, equal to $\hat{f}(u, v)$ on outcomes $u > \rho$ and satisfies (23) at ρ that has strictly lower variance than $\hat{f}$ on all vectors $z \in Z \cap V^*(\rho, v)$. This will show that $\hat{f}$ is not variance⁺-optimal on Z .

The estimator $\hat{f}$ violates (24), so either

$$\lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} < \inf_{z \in Z \cap V^*(\rho, v)} \lambda(\rho, z) \equiv L \quad (25)$$

or

$$\lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} > \sup_{z \in Z \cap V^*(\rho, v)} \lambda(\rho, z) \equiv U. \quad (26)$$

Violation (26), for a nonnegative unbiased $\hat{f}$, means that $M \equiv \int_{\rho}^1 \hat{f}(u, v) du < \underline{f}(u, v)$. Consider $z \in Z \cap V^*(\rho, v)$ and the z -optimal extension, $\hat{f}^{(z, \rho, M)}$ (see Theorem 2.1). Because the point (ρ, M) lies strictly below $\underline{f}^{(z)}$, the lower hull of both the point and $\underline{f}^{(z)}$ has a linear piece on some interval with right end point ρ . More precisely, $\hat{f}^{(z, \rho, M)}(u) \equiv \lambda(\rho, z, M)$ on $S(u, z)$ at some nonempty interval $u \in (\eta_z, \rho]$ so that at the point η_z , the lower bound is met, that is, $M + (\rho - \eta_z)\lambda(\rho, z, M) = \lim_{u \rightarrow \eta_z^+} \underline{f}(u, z)$. Therefore, all extensions (maintaining nonnegativity and unbiasedness) must satisfy

$$\int_{\eta_z}^{\rho} \hat{f}(u, z) du \leq \lim_{u \rightarrow \eta_z^+} \underline{f}(u, z) - M \quad (27)$$

$$= (\rho - \eta_z)\lambda(\rho, z, M) \leq (\rho - \eta_z)U.$$

From (26), for some $\epsilon > 0$, $\hat{f}$ has average value strictly higher than U on $S(u, v)$ for all u in intervals $(\eta, \rho]$ for $\eta \in [\rho - \epsilon, \rho)$. For each $z \in V^*(\rho, v)$ we define ζ_z as the maximum of $\rho - \epsilon$ and $\inf\{u | V^*(u, v) = V^*(\rho, v)\}$. From (1), $\zeta_z < \rho$. For each z , the higher estimate values on $S(u, z)$ for $u \in (\zeta_z, \rho]$ must be “compensated for” by lower values on $u \in (\eta_z, \zeta_z)$ (from nonnegativity we must have $\eta_z < \zeta_z$) so that (27) holds. By modifying the estimator to be equal to U for all outcomes $S(u, v)$ $u \in (\rho - \epsilon, \rho]$ and correspondingly increasing some estimate values that

are lower than U to U on $S(u, z)$ for $u \in (\eta_z, \zeta_z)$ we obtain an estimator with strictly lower variance than $\hat{f}$ for all $z \in Z \cap V^*(\rho, v)$ and same variance as $\hat{f}$ on all other vectors. Note we can perform the shift consistently across all branches of the tree-like partial order on outcomes.

Violation (25) means that for some $\epsilon > 0$, $\hat{f}$ has average value strictly lower than L on $S(u, v)$ for all intervals $u \in (\eta, \rho]$ for $\eta \in [\rho - \epsilon, \rho)$. For all z , the z -optimal extension $\hat{f}^{(z, \rho, M)}(u)$ has value $\lambda(\rho, z, M) \geq L$ at ρ and (from convexity of lower hull) values that are at least that on $u < \rho$. From unbiasedness, we must have for all $z \in Z \cap V^*(\rho, v)$, $\int_0^{\rho} \hat{f}(u, z) du = \int_0^{\rho} \hat{f}^{(z, \rho, M)}(u) du$. Therefore, values lower than L must be compensated for in $\hat{f}$ by values higher than L . We can modify the estimator such that it is equal to L for $S(u, v)$ for $u \in (\rho - \epsilon, \rho)$ and compensate for that by lowering values at lower u values $u < \zeta_z$ that are higher than L . The modified estimator has strictly lower variance than $\hat{f}$ for all $z \in Z \cap V^*(\rho, v)$ and same variance as $\hat{f}$ on all other vectors. $\square$

4. THE L ESTIMATOR

The estimator $\hat{f}^{(L)}$ satisfies (19a) with equalities, obtaining values that are minimum in the pointwise optimal range. Its values on outcomes consistent with data v only depend on the lower bound function on outcomes consistent with v . Geometrically, as visualized in Figure 2, the L estimate on an outcome $S(\rho, v)$ is exactly the slope value that if maintained for outcomes $S(u, v)$ ($u \in (0, \rho]$), would yield an expected estimate of $\underline{f}(S)$. We derive a convenient expres-




Figure 2: An example lower bound function $\underline{f}^{(v)}(u)$ with 3 steps and the respective cumulative L estimate $\int_u^1 \hat{f}^{(L)}(u, v) du$. The estimate $\hat{f}^{(L)}$ is the negated slope and in this case is also a step function with 3 steps.

sion for this estimator and show that it is 4-competitive and that it is the unique variance⁺ optimal monotone estimator. We also show it is order-optimal with respect to the natural order that prioritizes data vectors with lower $f(v)$.

The L estimator is the solution of the integral equation

$\forall \mathbf{v} \in \mathbf{V}, \forall \rho \in [0, 1]$:

$$\hat{f}^{(L)}(\rho, \mathbf{v}) = \frac{\underline{f}^{(\mathbf{v})}(\rho) - \int_{\rho}^1 \hat{f}^{(L)}(u, \mathbf{v}) du}{\rho} \quad (28)$$

We solve the integral equations to obtain an explicit form of the estimator $\hat{f}^{(L)}$.

LEMMA 4.1.

$$\hat{f}^{(L)}(\rho, \mathbf{v}) = \frac{\underline{f}^{(\mathbf{v})}(\rho)}{\rho} - \int_{\rho}^1 \frac{\underline{f}^{(\mathbf{v})}(u)}{u^2} du \quad (29)$$

$$= - \int_{\rho}^1 \frac{\frac{d\underline{f}^{(\mathbf{v})}(u)}{du}}{u} du \quad (30)$$

PROOF. We show that (29) is a solution of (28). To ease calculations, we use $h(x) \equiv \int \frac{\underline{f}^{(\mathbf{v})}(x)}{x^2} dx$. We have

$$\hat{f}^{(L)}(\rho, \mathbf{v}) = \frac{\underline{f}^{(\mathbf{v})}(\rho)}{\rho} - h(1) + h(\rho). \quad (31)$$

We need to show

$$\underline{f}^{(\mathbf{v})}(\rho) = \rho \hat{f}^{(L)}(\rho, \mathbf{v}) + \int_{\rho}^1 \hat{f}^{(L)}(x, \mathbf{v}) dx$$

Substituting (31) we get

$$\underline{f}^{(\mathbf{v})}(\rho) = \underline{f}^{(\mathbf{v})}(\rho) - \rho(h(1) - h(\rho)) + \int_{\rho}^1 \left(\frac{\underline{f}^{(\mathbf{v})}(x)}{x} - h(1) + h(x) \right) dx$$

$$\rho(h(1) - h(\rho)) = \int_{\rho}^1 \left(\frac{\underline{f}^{(\mathbf{v})}(x)}{x} + h(x) \right) dx - (1 - \rho)h(1)$$

Rearranging and canceling identical terms

$$h(1) - \rho h(\rho) = \int_{\rho}^1 \left(\frac{\underline{f}^{(\mathbf{v})}(x)}{x} - h(x) \right) dx \quad (32)$$

Using $(h(x)x)' = xh'(x) + h(x)$, and substituting $h'(x)x = \underline{f}^{(\mathbf{v})}(x)/x$ we obtain that $h(x)x = \int (\underline{f}^{(\mathbf{v})}(x)/x + h(x)) dx$. Substituting in (32) we get that we need to show

$$h(1) - \rho h(\rho) = h(x)x|_{\rho}^1 = h(1) - \rho h(\rho)$$

Lastly, the lower bound function $\underline{f}^{(\mathbf{v})}(u)$ is monotone on $(0, 1]$ and thus differentiable almost everywhere. Thus, $\frac{d\underline{f}^{(\mathbf{v})}(u)}{du}$ is defined almost everywhere. We get (30) from (29) using integration by parts. $\square$

We show a tight bound of 4 for the competitive ratio for $\hat{f}^{(L)}$, meaning that it is at most 4 for all functions f and for any $\epsilon > 0$, there exists a function f on which the ratio is no less than $4 - \epsilon$.

THEOREM 4.1.

$$\sup_{f, \mathbf{v} \mid \int_0^1 \hat{f}^{(\mathbf{v})}(u)^2 du < \infty} \frac{\int_0^1 \hat{f}^{(L)}(u, \mathbf{v})^2 du}{\int_0^1 \hat{f}^{(\mathbf{v})}(u)^2 du} = 4,$$

We first present a family of functions for which the supremum of this ratio is 4. Consider the domain $[0, 1]^2$, $\tau(u) = u$, and $f(\mathbf{v}) = \frac{1}{1-\alpha} - \frac{\max(\mathbf{v})^{1-\alpha}}{1-\alpha}$ for $\alpha < 0.5$. For the vector $\mathbf{v} = (0, 0)$ we have the following convex lower bound function

$$\underline{f}(u, (0, 0)) = \frac{1}{1-\alpha} - \frac{u^{1-\alpha}}{1-\alpha}.$$

Alternatively, the same lower bound function also corresponds to data $\mathbf{v} = (1, 0)$ and the function $g(\mathbf{v}) = \frac{1}{1-\alpha} - \frac{\text{RG}(\mathbf{v})^{1-\alpha}}{1-\alpha}$.

Being convex, this lower bound function is equal to its lower hull. Therefore, by taking its negated derivative, we get $\hat{f}^{(0,0)}(u) = 1/u^\alpha$. The function $\hat{f}^{(0,0)}$ is square integrable when $\alpha < 0.5$:

$$\int_0^1 \hat{f}^{(0,0)}(u)^2 du = \int_0^1 1/u^{2\alpha} du = \frac{1}{1-2\alpha}.$$

From (30), the L estimator on outcomes consistent with $\mathbf{v}$ is

$$\hat{f}^{(L)}(x, \mathbf{v}) = \int_x^1 \frac{1}{u^{1+\alpha}} = \frac{1}{\alpha} \left(\frac{1}{x^\alpha} - 1 \right).$$

Hence,

$$\begin{aligned} \int_0^1 \hat{f}^{(L)}(u, \mathbf{v})^2 du &= \frac{1}{\alpha^2} \int_0^1 \left(\frac{1}{u^{2\alpha}} - \frac{2}{u^\alpha} + 1 \right) du \\ &= \frac{1}{\alpha^2} \left(\frac{1}{1-2\alpha} - \frac{2}{1-\alpha} + 1 \right) = \frac{2}{(1-2\alpha)(1-\alpha)}. \end{aligned}$$

We obtain the ratio

$$\frac{\int_0^1 \hat{f}^{(L)}(u, \mathbf{v})^2 du}{\int_0^1 \hat{f}^{(\mathbf{v})}(u)^2 du} = \frac{2}{1-\alpha} \leq 4.$$

The ratio approaches 4 when $\alpha \rightarrow 0.5^-$.

We conclude the proof of Theorem 4.1 using the following lemma that shows that if $\hat{f}^{(\mathbf{v})}(u)$ is square integrable, that is, (8) holds, then $\hat{f}^{(L)}(u, \mathbf{v})$ is also square integrable and the ratio between these integrals is at most 4.

LEMMA 4.2.

$$\forall \mathbf{v}, \int_0^1 \hat{f}^{(\mathbf{v})}(u)^2 du < \infty \implies \frac{\int_0^1 \hat{f}^{(L)}(u, \mathbf{v})^2 du}{\int_0^1 \hat{f}^{(\mathbf{v})}(u)^2 du} \leq 4$$

PROOF. The function $\hat{f}^{(\mathbf{v})}$ only depends on the lower hull of the lower bound function $\underline{f}^{(\mathbf{v})}(u)$. The estimator $\hat{f}^{(L)}$ depends on the lower bound function $\underline{f}$ and can be different for different lower bound functions with the same lower hull. Fixing the lower hull, the variance of the L estimator is maximized for f such that $\underline{f}^{(\mathbf{v})} \equiv H_f^{(\mathbf{v})}$. It therefore suffices to consider convex $\underline{f}^{(\mathbf{v})}(u)$, that is, $\frac{d^2 \underline{f}^{(\mathbf{v})}(u)}{d^2 u} > 0$ for which we have

$$\hat{f}^{(\mathbf{v})}(u) = - \frac{d\underline{f}^{(\mathbf{v})}(u)}{du}.$$

Recall that $\hat{f}^{(\mathbf{v})}(u)$ is monotone non-increasing. From (30),

$$\hat{f}^{(L)}(\rho, \mathbf{v}) = - \int_{\rho}^1 \frac{\frac{d\underline{f}^{(\mathbf{v})}(u)}{du}}{u} du.$$

To establish our claim, it suffices to show that for all monotone non increasing functions g ,

$$\frac{\int_0^1 (\int_x^1 \frac{g(u)}{u} du)^2 dx}{\int_0^1 g(x)^2 dx} \leq 4 \quad (33)$$

Define $h(x) = \int_x^1 \frac{g(u)}{u} du$.

$$\begin{aligned} \int_\epsilon^1 h^2(x) dx &= \int_\epsilon^1 \int_x^1 2h(y)h'(y) dy dx \\ &= \int_\epsilon^1 \int_\epsilon^y 2h(y)h'(y) dx dy \\ &= 2 \int_\epsilon^1 h(y)h'(y) \int_\epsilon^y dx dy \\ &= 2 \int_\epsilon^1 h(y)h'(y)(y - \epsilon) dy \\ &= 2 \int_\epsilon^1 h(y) \frac{g(y)}{y} (y - \epsilon) dy \leq 2 \int_\epsilon^1 h(y)g(y) dy \\ &\leq 2 \sqrt{\int_\epsilon^1 h^2(y) dy} \sqrt{\int_\epsilon^1 g^2(y) dy} \end{aligned}$$

The last inequality is Cauchy-Schwartz. To obtain (33), we divide both sides by $\sqrt{\int_\epsilon^1 h^2(y) dy}$ and take the limit as ϵ goes to 0.

□

THEOREM 4.2. *The estimator $\hat{f}^{(L)}$ is monotone. Moreover, it is the unique variance⁺-optimal monotone estimator and dominates all monotone estimators.*

PROOF. Recall that an estimator $\hat{f}$ is monotone if and only if, for any data $\mathbf{v}$, the estimate $\hat{f}(\rho, \mathbf{v})$ is non-increasing with ρ . To show monotonicity of the L estimators, we rewrite (29) to obtain

$$\hat{f}^{(L)}(\rho, \mathbf{v}) = \underline{f}(\rho) + \int_\rho^1 \frac{\underline{f}(\rho) - \underline{f}(x)}{x^2} dx, \quad (34)$$

which is clearly non-increasing with ρ .

We now show that $\hat{f}^{(L)}$ dominates all monotone estimators (and hence is the unique variance⁺-optimal monotone estimator). By definition, a monotone estimator $\hat{f}$ must satisfy the inequalities $\forall \mathbf{v}, \forall \rho \in [0, 1]$:

$$\begin{aligned} \rho \hat{f}(\rho, \mathbf{v}) + \int_\rho^1 \hat{f}(u, \mathbf{v}) du &\leq \inf_{z \in V^*(\rho, \mathbf{v})} \int_0^1 \hat{f}(u, z) du = \\ \inf_{z \in V^*(\rho, \mathbf{v})} \underline{f}(z) &= \underline{f}(\rho, \mathbf{v}). \end{aligned} \quad (35)$$

Estimator $\hat{f}^{(L)}$ satisfies (35) with equalities. If there is a monotone estimator $\hat{f}$ with an inequality which is not tight, we can obtain a monotone estimator that strictly dominates $\hat{f}$ by decreasing the estimate for $u \leq \rho$ and increasing it for $u > \rho$. The variance decreases because we decrease the estimate on higher values and increase on lower values. □

Lastly, we show that $\hat{f}^{(L)}$ is order-based variance optimal with respect to the order $\prec$ which prioritizes vectors with lower $f(\mathbf{v})$:

THEOREM 4.3. *A $\prec^+$ -optimal estimator for f with respect to the partial order*

$$\mathbf{v} \prec \mathbf{v}' \iff f(\mathbf{v}) < f(\mathbf{v}')$$

must be equivalent to $\hat{f}^{(L)}$.

PROOF. We use our results of order-based optimality (Section 5). We can check that we obtain (28) using (42) and $\prec$ as defined in the statement of the Theorem. Thus, a $\prec^+$ -optimal solution must have this form. □

The L estimator may not be bounded (see Example 4). An estimator that is both bounded and competitive (but not necessarily in-range) is the J estimator [8]. For any $\epsilon > 0$, the estimator satisfying $\hat{f}(S) = \min\{(1 + \epsilon)\lambda_L(S), \lambda_U(S)\}$ is in-range, is bounded, if (9) holds, and is competitive, if (8) holds.

5. ORDER-BASED OPTIMALITY

We identify conditions on f and $\prec$ under which a $\prec^+$ -optimal estimator exists and specify this estimator as a solution of a set of equations. Our derivations of $\prec^+$ -optimal estimators follow the intuition to require the estimate on an outcome S to be $\mathbf{v}$ -optimal with respect to the $\prec$ -minimal consistent vector:

$$\forall S = S(\rho, \mathbf{v}), \hat{f}(S) = \lambda(\rho, \min_{\prec}(V^*(S))). \quad (36)$$

When $\prec$ is a total order and V is finite, $\min_{\prec}(V^*(S))$ is unique and (36) is well defined. Moreover, as long as f has a nonnegative unbiased estimator, a solution (36) always exists and is $\prec^+$ -optimal. We preview a simple construction of the solution: Process vectors in increasing $\prec$ order, iteratively building a partially defined nonnegative estimator. When processing $\mathbf{v}$, the estimator is already defined for $S(u, \mathbf{v})$ for $u \geq \rho_v$, for some $\rho_v \in (0, 1]$. We extend it to the outcomes $S(u, \mathbf{v})$ for $u \leq \rho_v$ using the $\mathbf{v}$ -optimal extension $\hat{f}^{(\mathbf{v}, \rho_v, M)}(u)$, where $M = \int_{\rho_v}^1 \hat{f}(u, \mathbf{v}) du$ (see Theorem 2.1).

We now formulate conditions that will allow us to establish $\prec^+$ -optimality of a solution of (36) in more general settings. These conditions always hold when $\prec$ is a total order and V is finite. Generally,

$$\min_{\prec}(V^*(S)) = \{z \in V^*(S) \mid \neg \exists \mathbf{w} \in V^*(S), \mathbf{w} \prec z\}$$


is a set and (36) is well defined when $\forall S$, this set is not empty and $\lambda(\rho, \min_{\prec}(V^*(S)))$ is unique, that is, the value $\lambda(\rho, z)$ is the same for all $\prec$ -minimal vectors $z \in \min_{\prec}(V^*(S))$. A sufficient condition for this is that

$$\begin{aligned} \forall \rho \forall \mathbf{v} \forall x \in (0, \underline{f}(\rho, \mathbf{v})] \forall z, \mathbf{w} \in \min_{\prec}(V^*(\rho, \mathbf{v})), \\ \inf_{\eta < \rho} \frac{\underline{f}(\eta, z) - x}{\rho - \eta} = \inf_{\eta < \rho} \frac{\underline{f}(\eta, \mathbf{w}) - x}{\rho - \eta} \end{aligned} \quad (37)$$

Example 4 L and U estimates for Example 3

We compute the L and U estimators for RG_{p+} for the sampling scheme and data in Example 3. For the two vectors $(0.6, 0.2)$ and $(0.6, 0)$, both the L and U estimates are 0 when $u \geq 0.6$, this is necessary from unbiasedness and nonnegativity because for these outcomes $\exists v \in V^*(S)$, $RG_{p+}(v) = 0$. Otherwise, the L estimate is $\hat{RG}_{p+}^{(L)}(S) = (v_1 - v'_2)^p / v'_2 - \int_{v'_2}^{v_1} \frac{(v_1 - x)^p}{x^2} dx$, where $v'_2 = u$ when $S = \{1\}$ and $v'_2 = v_2$ when $S = \{1, 2\}$. When $p \geq 1$, the U estimate is $\hat{RG}_{p+}^{(U)}(S) = p(v_1 - u)^{p-1}$ when $u \in (v_2, v_1]$ and 0 when $u \leq v_2 < v_1$. When $p \leq 1$ the U estimate is v_1^{p-1} when $u \in (v_2, v_1]$ and $\frac{(v_1 - v_2)^p - v_1^{p-1}(v_1 - v_2)}{v_2}$ when $u \leq v_2 < v_1$.

The figure also include the v -optimal estimates, discussed in Example 3. When $v_2 = 0$, the U estimates are v -optimal. The L estimate is not bounded when $v_2 = 0$ (but has bounded variance and is competitive).



In this case, the respective Equation (36) on $u \in (0, \rho]$ are the same for all $z \in \min_{\prec}(V^*(S))$ and thus so are the estimate values $\hat{f}(u, z)$.

We say that $Z \subset V$ is $\prec$ -bounded if

$$\forall v \in Z \exists z \in \min_{\prec}(Z), z \preceq v \quad (38)$$

That is, for all $z \in Z$, z is $\prec$ -minimal or is preceded by some vector that is $\prec$ -minimal in Z .

We say that an outcome S is $\prec$ -bounded if $V^*(S)$ is $\prec$ -bounded, that is,

$$\forall v \in V^*(S) \exists z \in \min_{\prec}(V^*(S)), z \preceq v \quad (39)$$

When all outcomes $S(u, v)$ are $\prec$ -bounded, we say that a set of vectors R represents v if any outcome consistent with v has a $\prec$ -minimal vector in R :

$$\forall u \in (0, 1], \exists z \in R, z \in \min_{\prec}(V^*(u, v)).$$

We now show that we can obtain a $\prec^+$ -optimal estimator if every vector v has a set of finite size that represents it. Example 5 (Appendix) walks through a derivation of $\prec^+$ -optimal estimators.

LEMMA 5.1. If f satisfies (7), (37), (39) and

$$\forall v, \min\{|R| \mid \forall u \in (0, 1], \exists z \in R, z \in \min_{\prec}(V^*(u, v))\} < \infty,$$

then a $\prec^+$ -optimal estimator exists and must be equivalent to a solution of (36).

PROOF. We provide an explicit construction of a $\prec^+$ -optimal estimator for f .

Fixing v , we select a finite set of representatives. We can map the representatives (or a subset of them) to distinct subintervals covering $(0, 1]$. The subintervals have the form

$(a_i, a_{i-1}]$ where $0 = a_n < \dots < a_1 < a_0 = 1$ such that a representative z that is minimal for $(a_i, a_{i-1}]$ is not minimal for $u \leq a_i$. Such mapping can always be obtained since from (1), each vector is consistent with an open interval of the form $(a, 1]$, and thus if $\prec$ -minimum at $V^*(u, v)$ (we must have $u > a$) it must be $\prec$ -minimum for $V^*(x, v)$ for $x \in (a, u]$. Thus, the region on which z is in $\min_{\prec} V^*(u, v)$ is open to the left. We can always choose a mapping such that the left boundary of this region corresponds to a_i .

Let $z^{(i)}$ ($i \in [n]$) be the representative mapped to outcomes $S(u, v)$ where $u \in (a_i, a_{i-1}]$. Since $V^*(u, v)$ is monotone non-decreasing with u , $i < j$ implies that $z^{(i)} \prec z^{(j)}$ or that they are incomparable in the partial order.

We construct a partially specified nonnegative estimator in steps, by solving (36) iteratively for the vectors $z^{(i)}$. Initially we invoke Theorem 2.1 to obtain estimate values for $S(u, z^{(1)})$ $u \in (0, 1]$ that minimize the variance for $z^{(1)}$. The result is a partially specified nonnegative estimator. In particular for v , the estimator is now specified for outcomes $S(u, v)$ where $u \in (a_1, 1]$. Any modification of this estimator on a subinterval of $(a_1, 1]$ with positive measure will strictly increase the variance for $z^{(1)}$ (or result in an estimator that can not be completed to a nonnegative unbiased one).

After step i , we have a partially specified nonnegative estimator that is specified for $S(u, v)$ for $u \in (a_i, 1]$. The estimator is fully specified for $z^{(j)}$ $j \leq i$ and is $\prec^+$ -optimal on these vectors in the sense that any other partially specified nonnegative estimator that is fully specified for $z^{(j)}$ $j \leq i$ and has strictly lower variance on some $z^{(j)}$ ($j \leq i$) must have strictly higher variance on some $z^{(h)}$ such that $h < j$.

We now invoke Theorem 2.1 with respect to the vector $z^{(i+1)}$. The estimator is partially specified for $S(u, z^{(i+1)})$

on $u > a_i$ and we obtain estimate values for the outcomes $S(u, z^{(i+1)})$ for $u \in (0, a_i]$ that constitute a partially specified nonnegative estimator with minimum variance for $z^{(i+1)}$. Note again that this completion is unique (up to equivalence). This extension now defines $S(u, v)$ for $u \in (a_{i+1}, 1]$.

Lastly, note that we must have $f(z^{(n)}) = f(v)$ because $f(z^{(n)}) < f(v)$ implies that (7) is violated for v whereas the reverse inequality implies that (7) is violated for $z^{(n)}$. Since at step n the estimator is specified for all outcomes $S(u, z^{(n)})$ and unbiased, it is unbiased for v .

The estimator is invariant to the choice of the representative sets R_v for $v \in V$ and also remains the same if we restrict $\prec$ so that it includes only relations between v and R_v .

We so far showed that there is a unique, up to equivalence, partially specified nonnegative estimator that is $\prec^+$ optimal with respect to a vector v and all vectors it depends on. Consider now all outcomes $S(u, v)$, for all u and v , arranged according to the containment order on $V^*(u, v)$ according to decreasing u values with branching points when $V^*(u, v)$ changes. If for two vectors v and z , the sets of outcomes $S(u, v), u \in (0, 1]$ and $S(u, z), u \in (0, 1]$ intersect, the intersection must be equal for $u > \rho$ for some $\rho < 1$. In this case the estimator values computed with respect to either z or v would be identical for $u \in (\rho, 1]$. Also note that partially specified nonnegative solutions on different branches are independent. Therefore, solutions with respect to different vectors v can be consistently combined to a fully specified estimator. $\square$

5.1 Continuous domains

The assumptions of Lemma 5.1 may break on continuous domains. Firstly, outcomes may not be $\prec$ -bounded and in particular, $\min_{\prec}(V^*(S))$ can be empty even when $V^*(S)$ is not, resulting in (36) not being well defined. Secondly, even if $\prec$ is a total order, minimum elements do not necessarily exist and thus (39) may not hold, and lastly, there may not be a finite set of representatives. To treat such domains, we utilize a notion of *convergence with respect to* $\prec$:

We define the $\prec$ -lim of a function h on a set of vectors $Z \subset V$:

$$\begin{aligned} \prec\text{-}\lim(h(\cdot), Z) = x &\iff \\ \forall v \in Z \forall \epsilon > 0 \exists w \preceq v, \forall z \preceq w, |h(z) - x| &\leq \epsilon \end{aligned} \quad (40)$$

The $\prec$ -lim may not exist but is unique if it does. Note that when Z is finite or more generally, $\prec$ -bounded, and $h(z)$ is unique for all $z \in \min_{\prec} Z$, then $\prec\text{-}\lim(h(\cdot), Z) = h(\min_{\prec} Z)$.

We define the $\prec$ -closure of z as the set containing z and all preceding vectors $\text{cl}_{\prec}(z) = \{v \in V | v \preceq z\}$.

We provide an alternative definition of the $\prec$ -lim using the notion of $\prec$ -closure.

$$\begin{aligned} \prec\text{-}\lim(h(\cdot), Z) = x &\quad (41) \\ \iff \inf_{v \in Z} \sup_{z \in \text{cl}_{\prec}(v) \cap Z} h(z) = \sup_{v \in Z} \inf_{z \in \text{cl}_{\prec}(v) \cap Z} h(z) = x \end{aligned}$$

We say that the lower bound function $\prec$ -converges on outcome $S = S(\rho, v)$ if $\prec\text{-}\lim(\underline{f}(\eta, \cdot), V^*(S))$ exists for all $\eta \in (0, \rho)$. When this holds, the $\prec$ -lim of the pointwise optimal values (15) over consistent vectors $V^*(S)$ exists for all $M = \int_{\rho}^1 \hat{f}(u, v) du \leq \underline{f}(\rho, v)$. We use the notation

$$\begin{aligned} \lambda_{\prec}(S, M) &= \prec\text{-}\lim(\lambda(\rho, \cdot, M), V^*(S)) \\ &= \inf_{0 \leq \eta < \rho} \frac{\prec\text{-}\lim(\underline{f}(\eta, \cdot), V^*(S)) - M}{\rho - \eta}. \end{aligned}$$

When the partially specified estimator $\hat{f}$ is clear from context, we omit the parameter M and use the notation

$$\begin{aligned} \lambda_{\prec}(S) &= \prec\text{-}\lim(\lambda(\rho, \cdot), V^*(S)) \\ &= \inf_{0 \leq \eta < \rho} \frac{\prec\text{-}\lim(\underline{f}(\eta, \cdot), V^*(S)) - \int_{\rho}^1 \hat{f}(u, v) du}{\rho - \eta}. \end{aligned}$$

We can finally propose a generalization of (36):

$$\forall S, \hat{f}(S) = \lambda_{\prec}(S) \quad (42)$$

which is well defined when the lower bound function $\prec$ -converges for all S :

$$\forall S = S(\rho, v), \forall \eta \leq \rho, \prec\text{-}\lim(\underline{f}(\eta, \cdot), V^*(S)) \text{ exists.} \quad (43)$$

Using the definition (41) of $\prec$ -convergence and (2) we obtain that an estimator is equivalent to (42) if and only if

$$\forall v \forall \rho \in (0, 1], \lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} = \lambda_{\prec}(\rho, v) \quad (44)$$

We show that equivalence to (42) is *necessary* for $\prec^+$ -optimality. To facilitate the proof, we express $\prec^+$ -optimality in terms of restricted variance⁺ optimality:

LEMMA 5.2. *An estimator is $\prec^+$ -optimal if and only if, for all $v \in V$, it is variance⁺-optimal with respect to $\text{cl}_{\prec}(v)$.*

PROOF. If there is v such that $\hat{f}$ is not variance⁺-optimal on $\text{cl}_{\prec}(v)$, there is an alternative estimator with strictly lower variance on some $z \in \text{cl}_{\prec}(v)$ and at most the variance on all $\text{cl}_{\prec}(v) \setminus \{z\}$. Since $\text{cl}_{\prec}(v)$ contains all vectors that precede z , the estimator $\hat{f}$ can not be $\prec^+$ -optimal. To establish the converse, assume an estimator $\hat{f}$ is variance⁺ optimal on $\text{cl}_{\prec}(v)$ for all v . Consider $z \in V$. Since $\hat{f}$ is variance⁺-optimal on $\text{cl}_{\prec}(z)$, there is no alternative estimator with strictly lower variance on z and at most the variance of $\hat{f}$ on all preceding vectors. Since this holds for all z , we obtain that $\hat{f}$ is $\prec^+$ -optimal. $\square$

LEMMA 5.3. *If f satisfies (7) and (43) then $\hat{f}$ is $\prec^+$ -optimal only if it satisfies (44).*

PROOF. Lemma 5.2 states that an estimator is $\prec^+$ -optimal if and only if $\forall v \in V$ it is variance⁺-optimal with respect

to $\text{cl}_{\prec}(w)$. Applying Lemma 3.1, the latter holds only if

$$\begin{aligned} \forall v \in V \forall \rho \in (0, 1] \quad (45) \\ \lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} &\geq \inf_{z \in \text{cl}_{\prec}(v) \cap V^*(\rho, v)} \lambda(\rho, z) \\ &\leq \sup_{z \in \text{cl}_{\prec}(v) \cap V^*(\rho, v)} \lambda(\rho, z) \end{aligned}$$

From definition, $S(\rho, z) \equiv S(\rho, v)$ for all vectors $z \in V^*(\rho, v)$. Moreover, for $z \in V^*(\rho, v)$ there is a nonempty interval $(\eta_z, \rho]$ such that $\forall u \in (\eta_z, \rho]$, $V^*(u, z) \equiv V^*(u, v)$. Therefore, for all $z \in V^*(\rho, v)$, the limits $\lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, z) du}{\rho - \eta}$ are the same. Therefore, (45) $\iff$

$$\begin{aligned} \forall v \in V \forall \rho \in (0, 1] \quad (46) \\ \lim_{\eta \rightarrow \rho^-} \frac{\int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} &\geq \sup_{w \in V^*(\rho, v)} \inf_{z \in \text{cl}_{\prec}(w) \cap V^*(\rho, v)} \lambda(\rho, z) \\ &\leq \inf_{w \in V^*(\rho, v)} \sup_{z \in \text{cl}_{\prec}(w) \cap V^*(\rho, v)} \lambda(\rho, z) \end{aligned}$$

□

We leave open the question of determining the most inclusive conditions on f and $\prec$ under which a $\prec^+$ -optimum exists, and thus the solution of (42) is $\prec^+$ -optimal. We show that any solution of (42) is unbiased and nonnegative when f has a nonnegative unbiased estimator.

LEMMA 5.4. *When f and $\prec$ satisfy (7) and (43), a solution $\hat{f}^{(\prec^+)}$ of (42) is unbiased and nonnegative.*

PROOF. From Lemma 3.1, since all values are in-range, the solution is unbiased and nonnegative. □

6. THE U ESTIMATOR

The estimator $\hat{f}^{(U)}$ satisfies (19b) with equality.

$$\forall S(\rho, v), \hat{f}(\rho, v) = \sup_{z \in V^*(S)} \inf_{0 \leq \eta < \rho} \frac{f(\eta, z) - \int_{\eta}^{\rho} \hat{f}(u, v) du}{\rho - \eta} \quad (47)$$

The U estimator is not always variance $^+$ -optimal. We do show, however, that under a natural condition, it is order-based optimal with respect to an order that prioritizes vectors with higher f values. The condition states that for all $S(\rho, v)$ and $\eta < \rho$, the supremum of the lower bound function $\underline{f}(\eta, z)$ over $z \in V^*(S)$ is attained (in the limiting sense) at vectors that maximize f on $V^*(S)$. Formally:

$$\forall \eta < \rho, \lim_{x \rightarrow \bar{f}(S)} \sup_{z \in V^*(S) | f(z) \geq x} \underline{f}(\eta, z) = \sup_{z \in V^*(S)} \underline{f}(\eta, z), \quad (48)$$

where $\bar{f}(S) = \sup_{z \in V^*(S)} f(z)$.

LEMMA 6.1. *If f satisfies (48), then the U estimator is $\prec^+$ -optimal with respect to the order $z \prec v \iff f(z) > f(v)$.*

PROOF. We can show that when (48) holds then (47) is the same as (42). □

The condition (48) is satisfied by RG_p and RG_{p+} . In this case, the conditions of Lemma 5.1 are also satisfied and thus the U estimator is $\prec^+$ optimal.

Conclusion

We argued for the value of customizing estimators to patterns in the data while retaining performance guarantees on a larger domain, taking an optimization approach to estimators derivation. We demonstrated our approach for coordinated shared-seed samples and believe we provided foundations for going further, designing automated tools to derive estimators according to specifications and for exploring customization for other sampling schemes, in particular, independent sampling.

Acknowledgement

The authors would like to thank Micha Sharir for his help in the proof of Lemma 4.2.

7. REFERENCES

- [1] K. R. W. Brewer, L. J. Early, and S. F. Joyce. Selecting several samples from a single population. *Australian Journal of Statistics*, 14(3):231–239, 1972.
- [2] A. Z. Broder. On the resemblance and containment of documents. In *Proceedings of the Compression and Complexity of Sequences*, pages 21–29. IEEE, 1997.
- [3] A. Z. Broder. Identifying and filtering near-duplicate documents. In *Proceedings of the 11th Annual Symposium on Combinatorial Pattern Matching*, volume 1848 of *LLNCS*, pages 1–10. Springer, 2000.
- [4] E. Cohen. Size-estimation framework with applications to transitive closure and reachability. *J. Comput. System Sci.*, 55:441–453, 1997.
- [5] E. Cohen and H. Kaplan. Leveraging discarded samples for tighter estimation of multiple-set aggregates. In *Proceedings of the ACM SIGMETRICS’09 Conference*, 2009.
- [6] E. Cohen and H. Kaplan. Get the most out of your sample: Optimal unbiased estimators using partial information. In *Proc. of the 2011 ACM Symp. on Principles of Database Systems (PODS 2011)*. ACM, 2011. full version: <http://arxiv.org/abs/1203.4903>.
- [7] E. Cohen and H. Kaplan. How to estimate change from samples. Technical Report cs.DS/1203.4903, arXiv, 2012.
- [8] E. Cohen and H. Kaplan. What you can do with coordinated samples. Technical Report cs.DB/1206.5637, arXiv, 2012.
- [9] E. Cohen, H. Kaplan, and S. Sen. Coordinated weighted sampling for estimating aggregates over multiple weight assignments. *Proceedings of the VLDB Endowment*, 2(1–2), 2009. full version: <http://arxiv.org/abs/0906.4560>.
- [10] D. G. Horvitz and D. J. Thompson. A generalization of sampling without replacement from a finite universe. *Journal of the American Statistical Association*, 47(260):663–685, 1952.
- [11] E. Ohlsson. Coordination of pps samples over time. In *The 2nd International Conference on Establishment Surveys*, pages 255–264. American Statistical Association, 2000.
- [12] B. Rosén. Asymptotic theory for order sampling. *J. Statistical Planning and Inference*, 62(2):135–158, 1997.
- [13] P. J. Saavedra. Fixed sample size pps approximations with a permanent random number. In *Proc. of the Section on Survey Research Methods, Alexandria VA*, pages 697–700. American Statistical Association, 1995.

APPENDIX

Example 5 Walk-through derivation of $\prec^+$ -optimal estimators

We derive $\prec^+$ -optimal RG_{1+} estimators over the discrete domain $\mathbf{V} = \{0, 1, 2, 3\}^2$. Assuming same sampling scheme on both entries, there are 3 threshold values of interest, where π_i $i \in [3]$ is such that entry of value i is sampled if and only if $u \leq \pi_i$. We have $\pi_1 < \pi_2 < \pi_3$.

The lower bounds $\text{RG}_{1+}^{(v)}$ are step functions with steps at $u = \pi_i$. The table shows $\text{RG}_{1+}^{(v)}(u)$ for all u and v .

$\text{RG}_{1+}^{(v)}$	(1, 0)	(2, 1)	(2, 0)	(3, 2)	(3, 1)	(3, 0)
(0, π_1]	1	1	2	1	2	3
(π_1 , π_2]	0	1	1	1	2	2
(π_2 , π_3]	0	0	0	1	1	1
(π_3 , 1]	0	0	0	0	0	0

The v -optimal estimate, $\hat{\text{RG}}_{1+}^{(v)}(u)$ is the negated slope at u of the lower hull of $\text{RG}_{1+}^{(v)}$. The lower hull of each step function is piecewise linear with breakpoints at a subset of π_i , and thus, the v -optimal estimates are constant on each segment $(\pi_{i-1}, \pi_i]$. The table shows the estimates for all v and u . The notation $\downarrow$ refers to value in same column and row below and $\Downarrow$ to value two rows below.

$\hat{\text{RG}}_{1+}^{(v)}$	(1, 0)	(2, 1)	(2, 0)	(3, 2)	(3, 1)	(3, 0)
(0, π_1]	$\frac{1}{\pi_1}$	$\frac{1}{\pi_2}$	$\frac{2 - (\pi_2 - \pi_1)\downarrow}{\pi_1}$	$\frac{1}{\pi_3}$	$\frac{2 - \Downarrow}{\pi_2}$	$\frac{3 - \Downarrow(\pi_3 - \pi_2) - \Downarrow(\pi_2 - \pi_1)}{\pi_1}$
(π_1 , π_2]	0	$\frac{1}{\pi_2}$	$\min\{\frac{2}{\pi_2}, \frac{1}{\pi_2 - \pi_1}\}$	$\frac{1}{\pi_3}$	$\frac{2 - \downarrow}{\pi_2}$	$\min\{\frac{3 - \downarrow(\pi_3 - \pi_2)}{\pi_2}, \frac{2 - \downarrow(\pi_3 - \pi_2)}{\pi_2 - \pi_1}\}$
(π_2 , π_3]	0	0	0	$\frac{1}{\pi_3}$	$\min\{\frac{2}{\pi_3}, \frac{1}{\pi_3 - \pi_2}\}$	$\min\{\frac{3}{\pi_3}, \frac{1}{\pi_3 - \pi_2}\}$

The order $(2, 1) \prec (2, 0)$ and $(3, 2) \prec (3, 1) \prec (3, 0)$ yields the L estimator, which is v -optimal for $(1, 0)$, $(2, 1)$, and $(3, 2)$. The order $(2, 0) \prec (2, 1)$ and $(3, 0) \prec (3, 1) \prec (3, 2)$ yields the U estimator which is v -optimal for $(1, 0)$, $(2, 0)$, and $(3, 0)$.

To specify an estimator, we need to specify it on all possible outcomes, where each distinct outcome is uniquely determined by the sets $V^*(S)$. The 8 possible outcomes (we exclude those consistent with vectors with $\text{RG}_{1+}(v) = 0$ on which the estimate must be 0) are $(1, 0)$, $(2, \leq 1)$, $(2, 1)$, $(3, \leq 2)$, $(3, 2)$, $(3, \leq 1)$, $(3, 1)$, and $(3, 0)$.

We show how we construct the $\prec^+$ -optimal estimator for $\prec$ which prioritizes vectors with difference of 2: $(3, 1) \prec (3, 2) \prec (3, 0)$ and $(2, 0) \prec (2, 1)$.

The estimator is v -optimal for $(3, 1)$, $(2, 0)$, and $(1, 0)$. This determines the estimates $\hat{\text{RG}}_{1+}^{(\prec)}$ on all outcomes consistent with these vectors: The value on outcome $(1, 0)$ is $\hat{\text{RG}}^{((1,0))}((0, \pi_1])$, the values on outcomes $(2, \leq 1)$ and $(2, 0)$ are according to $\hat{\text{RG}}^{(2,0)}$ on $(\pi_1, \pi_2]$ and $(0, \pi_1]$, respectively, and value on the outcomes $(3, \leq 2)$, $(3, \leq 1)$ and $(3, 1)$ is according to $\hat{\text{RG}}^{(3,1)}$ on $(\pi_2, \pi_3]$ and $(\pi_1, \pi_2]$. These values are provided in the table above. The remaining outcomes are $(3, 0)$, $(3, 2)$, and $(2, 1)$. We need to specify the estimator so that it is unbiased on these vectors, given the existing specification. We have

$$\begin{aligned}\hat{\text{RG}}_{1+}^{(\prec)}(2, 1) &= \frac{1 - (\pi_2 - \pi_1)\hat{\text{RG}}_{1+}^{(\prec)}(2, \leq 1)}{\pi_1} \\ \hat{\text{RG}}_{1+}^{(\prec)}(3, 0) &= \frac{3 - (\pi_3 - \pi_2)\hat{\text{RG}}_{1+}^{(\prec)}(3, \leq 2) - (\pi_2 - \pi_1)\hat{\text{RG}}_{1+}^{(\prec)}(3, \leq 1)}{\pi_1} \\ \hat{\text{RG}}_{1+}^{(\prec)}(3, 2) &= \frac{2 - (\pi_3 - \pi_2)\hat{\text{RG}}_{1+}^{(\prec)}(3, \leq 2)}{\pi_1}.\end{aligned}$$

Observe that to apply these estimators, we do not have to precompute the estimator on all possible outcomes. An estimate only depends on values of the estimate on all less informative outcomes. In a discrete domain as in this example, it is the number of breakpoints larger than the seed u (which is at most the size of the domain).
